

NASA
Technical
Paper
3350

August 1993

**Semiempirical Fragmentation
Models on Galactic Cosmic
Ray Transport Calculations
With Hydrogen Target**

Judy L. Shinn,
John W. Wilson,
and Francis F. Badavi


(NASA-TP-3350) SEMIEMPIRICAL
FRAGMENTATION MODELS ON GALACTIC
COSMIC RAY TRANSPORT CALCULATIONS
WITH HYDROGEN TARGET (NASA) 18 p

N94-13263

Unclass

H1/93 0182494

NASA



1993

**Semiempirical Fragmentation
Models on Galactic Cosmic
Ray Transport Calculations
With Hydrogen Target**

Judy L. Shinn
and John W. Wilson
Langley Research Center
Hampton, Virginia

Francis F. Badavi
Christopher Newport University
Newport News, Virginia



National Aeronautics and
Space Administration
Office of Management
Scientific and Technical
Information Program

Abstract

Nuclear fragmentation cross sections of Silberberg and Tsao that are more accurate for a hydrogen target have been implemented in the data base to replace those of Rudstam for a galactic cosmic ray transport code (HZETRN). Sample calculations have been made for the transported galactic cosmic ray flux through a liquid hydrogen shield at solar minimum condition to determine the effect of such a change. The transported flux based on the Silberberg-Tsao semiempirical formalism contains fewer high-LET (linear energy transfer) components but more low-LET components than the results based on Rudstam's formalism; and this disparity deepens as the shield thickness increases. A comparison of the results obtained from using both energy-dependent and energy-independent cross sections of Silberberg and Tsao indicates that the energy-independent assumption results in an underestimation of high-LET flux above 100 keV/ μm by approximately 40 percent for a 15-g/cm² thickness of liquid hydrogen. Similar results were obtained in a previous study (NASA TP-3243) when both energy-dependent and energy-independent cross sections of Rudstam were considered. Nonetheless, the present study found that an energy-independent calculation would be best accomplished by using Rudstam's cross sections as done in the past for various engineering applications.

Introduction

Estimates of galactic cosmic ray exposure and the required shielding for astronauts during their interplanetary mission rely on radiation transport codes that accurately describe the interactions and propagation of these radiation fields throughout the bulk medium (ref. 1). HZETRN (ref. 2), which is a state-of-the-art galactic cosmic ray transport code developed at the Langley Research Center, contains detailed descriptions of physical processes such as energy loss, nuclear absorption, and fragmentation cross sections of projectile ions. However, the uncertainties of nuclear fragmentation cross sections existing in the code remain fairly large because of the lack of reliable experimental data (ref. 3). This condition is especially true for collisions of heavy ions on heavy ions for which the results of NUCFRG (ref. 4), based on a semiempirical abrasion-ablation model, are used as inputs to HZETRN. (Note that an improved version of a heavy ion fragmentation model will be available for use as inputs in the immediate future.) For collisions of heavy ions on protons, Rudstam's semiempirical parameterization (ref. 5) has been used in the existing HZETRN.

Rudstam and Metropolis et al. (refs. 5 and 6, respectively) were the pioneers who systematized high-energy cross section measurements into a useful analytical relationship describing nuclear reactions. These analytical relationships have been re-

vised and improved by many new researchers as new cross-section data become available. The most comprehensive set of semiempirical estimates of cross sections for a hydrogen target is perhaps due to Silberberg and Tsao (refs. 7 and 8), and to the most recent work of Webber, Kish, and Schrier (ref. 9). In the present study, the formulation of Silberberg and Tsao for a hydrogen target will replace that of Rudstam in HZETRN, and the effect of such a replacement will be examined for estimates of galactic cosmic ray exposure using liquid hydrogen as a shield.

Transport Methods

As energetic ions traverse through bulk matter, they lose energy through their interaction with atomic electrons along their trajectories. On occasion, they interact violently with nuclei of the matter and thus produce ion fragments or secondary nucleons moving in the forward direction and low-energy fragments of the struck target nucleus. The transport problem for the short-range target fragments can be easily solved in closed form in terms of collision density (ref. 10) and treated separately. Hence, the projectile-ion fragment or secondary nucleon transport is the remaining problem of interest. In the previous work (ref. 10), the projectile-ion fragments and secondary nucleons were treated as if all went straight ahead (ref. 11). The straight-ahead approximation is found to be quite accurate for the nearly isotropic cosmic ray fluence (ref. 10).

Because of the long range of the coulomb force and the large percentage of the material volume being occupied by electrons, the electron interactions can, to a good approximation, be treated as a continuous slowing-down process over any finite path length. Although the energy lost by an ion over some fixed path length fluctuates about a mean value, this fluctuation amounts to no more than a few percent (refs. 12-15) and is of no importance in the study of space radiation. In the following, a continuous slowing-down theory will be assumed, and the relevant quantity is the average loss per unit of path length.

With the straight-ahead approximation and the target secondary fragments neglected (refs. 10 and 11), the transport equation may be written as

$$\left[\frac{\partial}{\partial x} - \nu_j \frac{\partial}{\partial E} S(E) + \sigma_j(E) \right] \phi_j(x, E) = \sum_k \int_E^\infty f_{jk}(E, E') \phi_k(x, E') dE' \quad (1)$$

where $\phi_j(x, E)$ is the flux of ions of type j with atomic mass A_j and charge Z_j at x moving along the x -axis at energy E (in units of MeV/amu), σ_j is the corresponding macroscopic nuclear absorption cross section, $S(E)$ is the stopping power (continuous slowing-down approximation) of the protons, $f_{jk}(E, E')$ is a differential energy cross section for production of ion j in collision by ion k , and ν_j is the range scaling parameter that is defined as

$$\nu_j = \frac{Z_j^2}{A_j} \quad (2)$$

The solution to equation (1) is found to be subject to the boundary condition at $x = 0$ (that is, $\phi_j(0, E)$), which is the incident beam spectrum.

By transforming the heavy ion transport equation to an integral along the characteristic curve of that particular ion (ref. 10) and using the perturbation theory (ref. 16), the solution to equation (1) is given as a stepping procedure with step size h in the x -direction (ref. 17). Thus,

$$\begin{aligned} \psi_j(x + h, r) &\approx \exp[-\zeta_j(r, h)] \psi_j(x, r + \nu_j h) \\ &+ \sum_k \int_0^h \int_r^\infty \exp[-\zeta_j(r, z) - \zeta_k(r', h - z)] \bar{f}_{jk}(r + \nu_j z, r') \\ &\times \psi_k[x, r' + \nu_k(h - z)] dr' dz \end{aligned} \quad (3)$$

where

$$\psi_j(x, r_j) = S(E) \phi_j(x, E)$$

r_j is the residual range of ion j given by

$$r_j(E) = \int_0^E \frac{dE'}{S_j(E')}$$

and the exponential is the integrating factor with

$$\zeta_j(r, t) = \int_0^t \sigma_j(r + \nu_j t') dt'$$

Currently, we assume for $Z_j > 1$ and $k > j$ that

$$\bar{f}_{jk}(r, r') = f_{jk}(r') \delta(r - r') \quad (4)$$

Using equation (4), equation (3) now becomes

$$\begin{aligned}\psi_j(x+h, r) &\approx \exp[-\zeta_j(r, h)] \psi_j(x, r + \nu_j h) \\ &+ \sum_k \int_0^h dz \exp[-\zeta_j(r, z) - \zeta_k(r', h-z)] \sigma_{jk}(r') \\ &\times \psi_k[x, r' + \nu_k(h-z)]\end{aligned}\quad (5)$$

with $r' = r + \nu_j z$. Equation (5) is further approximated as

$$\begin{aligned}\psi_j(x+h, r) &\approx \exp[-\zeta_j(r, h)] \psi_j(x, r + \nu_j h) \\ &+ \sum_k \int_0^h dz \exp[-\zeta_j(r, z) - \zeta_k(r, h-z)] \sigma_{jk}(r) \psi_k[x, r + \nu_j z + \nu_k(h-z)] \\ &\approx \exp[-\sigma_j(r) h] \psi_j(x, r + \nu_j h) \\ &+ \sum_k \sigma_{jk}(r) \left\{ \frac{\exp[-\sigma_j(r) h] - \exp[-\sigma_k(r) h]}{\sigma_k(r) - \sigma_j(r)} \right\} \psi_k(x, r + \nu_j h) \\ &+ O[(\nu_k - \nu_j)h]\end{aligned}\quad (6)$$

Equation (6) is the stepping formalism with energy-dependent cross sections for $k > {}^4\text{He}$. The corresponding stepping formalism for nucleons has been discussed in detail in references 8–19. These stepping formalisms are then used to march the solution from the surface boundary to the desired shield thickness.

Nuclear Fragmentation Data Base

Even though the accuracy of the experimental data may improve for specific reactions, a reasonable means of representing data in computational procedures for cosmic ray transport calculation is still a challenge. For HZETRN and other radiation transport codes developed at Langley (refs. 18 and 20), a point representation of the data is avoided because large multidimensional arrays will eventually rival computer storage. Instead, various semi-empirical methods suitable for certain target or fragment groups are put together in generating the data base. These semiempirical methods, which are built on available experimental data and some theoretical base describing approximately the systematic variation of reaction cross sections, offer the possibility of implementing any additional necessary correction factors or adjusting some existing parameters.

When a nucleus is collided by high-energy nucleons, some individual nuclear constituents are ejected by direct knockout (ref. 21). The remaining nuclear structure is left in an excited state which seeks an equilibrium minimum-energy configuration through particle emission (ref. 5). This state is the basis of Rudstam's formulation for the systematics of spallation products produced in such collisions. In his formalism, the distribution of resultant isotopes that are related to the statistical nature of the evaporation process (ref. 22) is assumed to be Gaussian centered

at the nuclear stability line, and the total change in nuclear mass and the dependence on the incident projectile energy are treated empirically. Following Rudstam's work, Silberberg and Tsao (refs. 7 and 8) later added more corrective factors to the formalism as more experimental data became available. In addition, they added a scaling factor that relates the fragmentation cross section produced on a hydrogen target to that on the heavier target.

Samples of fragmentation cross section σ_{jk} used as inputs to HZETRN are shown as a function of projectile-ion charge Z_k and fragment-ion charge Z_j in figures 1 and 2 for a liquid hydrogen target and in figures 3 and 4 for a water target at various energies of the projectiles. Figures 1 and 3 reflect the results produced by using Rudstam's formalism for the fragments with $Z_j > 2$, whereas figures 2 and 4 reflect those by using the Silberberg-Tsao formalism. For nucleon and helium fragments, the cross sections were obtained from the results of Bertini et al. (refs. 23 and 24). (Note that the cross section shown in the figures has been reduced, for convenience, by a factor of 10 for nucleon fragments and a factor of 4 for helium fragments.) Because the scaling factor in the formalism of Silberberg and Tsao does not give adequate results for a target heavier than hydrogen (ref. 25), the semiempirical abrasion-ablation model (ref. 4) is used in obtaining the cross

sections contributed by the oxygen ions in the water target.

For all the cross sections shown in figures 1–4 at three separate projectile energies (150, 600, and 2400 MeV/amu), the peak of the fragments spectra (at constant Z_k) is generally higher as predicted by Rudstam than that predicted by Silberberg and Tsao. Conversely, the spread from the peak to the fragments with lower Z is lower for Rudstam, whereas the zig-zag shape over even and odd Z charges is more pronounced for Silberberg and Tsao. At energies below 150 MeV/amu, some results (not shown) given by Rudstam's formalism are erroneous; thus, for the transport calculations presented herein, the values of cross section at 150 MeV/amu have been extrapolated to below 150 MeV for projectile ions with $Z > 20$, as has been done in reference 17.

Results

By using Rudstam's formalism, several studies were made in the past that involved galactic cosmic ray (GCR) transport calculations through a shield that contained hydrogen atoms, such as water (tissue equivalent) (refs. 26 and 27) or liquid hydrogen (ref. 28). In these calculations, the cross sections were assumed to be energy independent and set equal to the asymptotic values at high energy. Recently, improvements were made to HZETRN by removing the energy-independent assumption; and the effects of such changes to the existing results were examined in terms of LET spectra that showed a substantial enhancement of high-LET components (ref. 17). In the present study, Rudstam's formalism was replaced by that of Silberberg and Tsao, and the effect of that replacement was then examined by comparing the GCR exposure levels behind the liquid hydrogen shield at the solar minimum condition given by the CREME model of reference 29.

The high-LET radiation components are usually degraded to lower LET components as a result of nuclear interactions between projectile and target nuclei, and such processes become more significant as the particles penetrate farther into the shield medium. This degradation is illustrated in figure 5 in which the annual differential dose and dose equivalent are plotted as a function of LET (or L) for 2-, 5-, and 15 g/cm² thicknesses of the shield. The spikes seen in the figures correspond to the zero slope of stopping power ($dS/dE = 0$) at minimum ionization (ref. 30) of each ion, with protons starting at the lowest LET followed by increasing Z for the increasing L . Because the LET coordinate is plotted on a logarithm scale, the differential dose $L dN/dL$ is converted to $L^2 dN/dL$ (where $N = \int \phi dE$) so that the

area under the curve is linearly proportional to the total dose. Similarly, the differential dose equivalent is plotted as $QL^2 dN/dL$, where the quality Q is a function of L . The magnification in dose equivalent at the high-LET region is a result of the ICRP 60 quality factor (ref. 31).

The differences between the spectra for the fragmentation cross section by Rudstam's formalism (fig. 1) and those by Silberberg and Tsao (fig. 2) as discussed earlier are reflected in the calculated LET spectra for the transported flux through the liquid hydrogen shield. In figure 6(a), the ratio of the LET spectra for the differential flux calculated by using Silberberg and Tsao relative to that by using Rudstam is displayed for the three thicknesses of the shield. The Silberberg-Tsao model produces more lower Z fragments and fewer higher Z fragments, and therefore more lower LET components and fewer higher LET components according to the spikes identified for each ion in figure 5. As a result, the ratio (fig. 6(a)) is higher at LET values near the 100-MeV/cm region and lower at higher values of LET. These deviations from unity become more pronounced as the shield thickness increases. A similar comparison of integral flux LET spectra is shown in figure 6(b) in which the ratio at the lowest LET is almost identical to unity; this indicates that the total flux does not change appreciably because of differences in the fragmentation model.

In the previous study (ref. 17), the removal of the energy-independent assumption was important in the risk assessment of GCR exposure. Using the liquid hydrogen shield as an example, the LET components above 100 keV/ μ m (or 1000 MeV/cm), which could contribute some biological-risk orders of magnitude higher than the lower components, were shown to increase by 40 percent at a 15-g/cm² thickness because of the removal of the assumption. This conclusion was based on the use of Rudstam's formalism. Similar results were obtained in figure 7 with the data base given by Silberberg and Tsao. Note in figure 5 that the region near 10 000 keV/ μ m was not as critical because of the diminishing flux in this region.

A comparison is made with the transport calculations using the energy-dependent Silberberg-Tsao data base and the energy-independent data base from Rudstam to serve as a reference for the earlier GCR exposure studies made with the old data base and the assumption of an energy-independent cross section which is taken to be the asymptotic value at 2 GeV/amu. The ratio of the two calculations in LET spectra, for both differential and integral flux shown in figures 8(a) and 8(b), respectively, displays

almost an oscillatory behavior over a wide range of LET. Thus, some of the differences are probably cancelled out. One arrives at the interesting observation that an energy-independent calculation using the Rudstam cross sections yields results similar to those of an energy-dependent calculation using the Silberberg-Tsao values.

Concluding Remarks


Nuclear fragmentation cross sections of Silberberg and Tsao that are more accurate for a hydrogen target have been placed in the data base for calculations of galactic cosmic ray (GCR) transport. When compared with the old data base of Rudstam, the Silberberg-Tsao model produces fewer higher charge fragments but more lower charge fragments. Sample calculations of GCR transport with a liquid hydrogen shield reflect such differences of cross sections in that the transported flux based on the Silberberg-Tsao model contains fewer high-LET (linear energy transfer) components but more lower LET components. This disparity deepens as the shield thickness increases. When the Silberberg-Tsao cross sections are assumed to be energy independent in the data base, the comparative results indicate an underestimation of the LET components above $100 \text{ keV}/\mu\text{m}$ by approximately 40 percent for a $15\text{-g}/\text{cm}^2$ thickness of liquid hydrogen caused by the assumption. Moreover, the present study found that an energy-independent calculation would be best accomplished by using Rudstam's cross sections as done in the past.

NASA Langley Research Center
Hampton, VA 23681-0001
June 8, 1993


References

1. Wilson, John W.; Townsend, Lawrence W.; Schimmerling, Walter; Khandelwal, Govind S.; Khan, Ferdous; Nealy, John E.; Cucinotta, Francis A.; Simonsen, Lisa C.; Shinn, Judy L.; and Norbury, John W.: *Transport Methods and Interactions for Space Radiations*. NASA RP-1257, 1991.
2. Wilson, John W.; Chun, Sang Y.; Badavi, Forooz F.; Townsend, Lawrence W.; and Lamkin, Stanley L.: *HZETRN: A Heavy Ion-Nucleon Transport Code for Space Radiations*. NASA TP-3146, 1991.
3. Townsend, Lawrence W.; Cucinotta, Francis A.; and Wilson, John W.: HZE Reactions and Data-Base Development. Paper presented at *NATO Advanced Study Institute on Biological Effects and Physics of Solar and Galactic Cosmic Radiation* (Amerção de Pera, Portugal), Oct. 12-23, 1991.
4. Wilson, John W.; Townsend, Lawrence W.; and Badavi, F. F.: A Semiempirical Nuclear Fragmentation Model. *Nucl. Instrum. & Methods Phys. Res.*, vol. B18, no. 3, Feb. 1987, pp. 225-231.
5. Rudstam, G.: Systematics of Spallation Yields. *Zeitschrift für Naturforschung*, vol. 21a, no. 7, July 1966, pp. 1027-1041.
6. Metropolis, N.; Bivins, R.; Storm, M.; Turkevich, Anthony; Miller, J. M.; and Friedlander, G.: Monte Carlo Calculations on Intranuclear Cascades. I. Low-Energy Studies. *Phys. Review*, second ser., vol. 110, no. 1, Apr. 1, 1958, pp. 185-203.
7. Silberberg, R.; and Tsao, C. H.: Partial Cross-Sections in High-Energy Nuclear Reactions, and Astrophysical Applications. I. Targets With $Z \leq 28$. *Astrophys. J. Suppl. Ser.*, no. 220(I), vol. 25, 1973, pp. 315-333.
8. Silberberg, R.; Tsao, C. H.; and Shapiro, M. M.: Semiempirical Cross Sections, and Applications to Nuclear Interactions of Cosmic Rays. *Spallation Nuclear Reactions and Their Applications*, B. S. P. Shen and M. Merker, eds., D. Reidel Publ. Co., c.1976, pp. 49-81.
9. Webber, W. R.; Kish, J. C.; and Schrier, D. A.: Formula for Calculating Partial Cross Sections for Nuclear Reactions of Nuclei With $E \gtrsim 200 \text{ MeV}/\text{Nucleon}$ in Hydrogen Targets. *Phys. Review C*, vol. 41, no. 2, Feb. 1990, pp. 566-571.
10. Wilson, John W.: *Analysis of the Theory of High-Energy Ion Transport*. NASA TN D-8381, 1977.
11. Wilson, John W.: *Heavy Ion Transport in the Straight Ahead Approximation*. NASA TP-2178, 1983.
12. Janni, Joseph F.: Proton Range-Energy Tables, 1 keV-10 GeV—Energy Loss, Range, Path Length, Time-of-Flight, Straggling, Multiple Scattering, and Nuclear Interaction Probability. Part 1. For 63 Compounds. *At. Data & Nucl. Data Tables*, vol. 27, nos. 2/3, Mar./May 1982, pp. 147-339.
13. Janni, Joseph F.: Proton Range-Energy Tables, 1 keV-10 GeV—Energy Loss, Range, Path Length, Time-of-Flight, Straggling, Multiple Scattering, and Nuclear Interaction Probability. Part 2. For Elements $1 \leq Z \leq 92$. *At. Data & Nucl. Data Tables*, vol. 27, nos. 4/5, July/Sept. 1982, pp. 341-529.
14. Schimmerling, Walter; Rapkin, Marwin; Wong, Mervyn; and Howard, Jerry: The Propagation of Relativistic Heavy Ions in Multielement Beam Lines. *Med. Phys.*, vol. 13, no. 2, Mar./Apr. 1986, pp. 217-228.
15. Alsmiller, R. G., Jr.; Barish, J.; and Scott, W. W.: The Effects of Multiple Coulomb Scattering and Range Straggling in Shielding Against Solar-Flare Protons. *Nucl. Sci. & Eng.*, vol. 35, no. 3, Mar. 1969, pp. 405-406.
16. Wilson, John W.; and Lamkin, Stanley L.: Perturbation Theory for Charged-Particle Transport in One Dimension. *Nucl. Sci. & Eng.*, vol. 57, no. 4, Aug. 1975, pp. 292-299.


17. Shinn, Judy L.; John, Sarah; Tripathi, Ram K.; Wilson, John W.; Townsend, Lawrence W.; and Norbury, John W.: *Fully Energy-Dependent HZETRN (A Galactic Cosmic-Ray Transport Code)*. NASA TP-3243, 1992.
18. Wilson, John W.; Townsend, Lawrence W.; Nealy, John E.; Chun, Sang Y.; Hong, B. S.; Buck, Warren W.; Lamkin, S. L.; Ganapol, Barry D.; Khan, Ferdous; and Cucinotta, Francis A.: *BRYNTRN: A Baryon Transport Model*. NASA TP-2887, 1989.
19. Shinn, Judy L.; Wilson, John W.; Weyland, Mark; and Cucinotta, Francis A.: *Improvements in Computational Accuracy of BRYNTRN (A Baryon Transport Code)*. NASA TP-3093, 1991.
20. Wilson, John W.; Lamkin, Stanley L.; Farhat, Hamidullah; Ganapol, Barry D.; and Townsend, Lawrence W.: *A Hierarchy of Transport Approximations for High Energy Heavy (HZE) Ions*. NASA TM-4118, 1989.
21. Serber, R.: Nuclear Reactions at High Energies. *Phys. Review*, vol. 72, no. 11, Dec. 1, 1947, pp. 1114-1115.
22. Dostrovsky, I.; Rabinowitz, P.; and Bivins, R.: Monte Carlo Calculations of High-Energy Nuclear Interactions. I. Systematics of Nuclear Evaporation. *Phys. Review*, vol. 111, no. 6, Sept. 15, 1958, pp. 1659-1676.
23. Bertini, Hugo W.; Guthrie, Miriam P.; and Culkowski, Arline H.: *Nonelastic Interactions of Nucleons and π -Mesons With Complex Nuclei at Energies Below 3 GeV*. CRNL-TM-3148, U.S. Atomic Energy Commission, Mar. 28, 1972.
24. *MECC-7 Intranuclear Cascade Code, 500-MeV Protons on 0-16*. I4C Analysis Codes (Programmed for H. W. Bertini). Available from Radiation Shielding Information Center, Oak Ridge National Lab., 1968.
25. Wilson, John W.; Townsend, Lawrence W.; and Khan, Ferdous: Evaluation of Highly Ionizing Components in High-Energy Nucleon Radiation Fields. *Health Phys.*, vol. 57, no. 5, Nov. 1989, pp. 717-724.
26. Shinn, Judy L.; Wilson, John W.; and Nealy, John E.: *Reliability of Equivalent Sphere Model in Blood-Forming Organ Dose Estimation*. NASA TM-4178, 1990.
27. Cucinotta, Francis A.; Atwell, William; Weyland, Mark; Hardy, Alva C.; Wilson, John W.; Townsend, Lawrence W.; Shinn, Judy L.; and Katz, Robert: *Radiation Risk Predictions for Space Station Freedom Orbits*. NASA TP-3098, 1991.
28. Townsend, Lawrence W.; Nealy, John E.; Wilson, John W.; and Simonsen, Lisa C.: *Estimates of Galactic Cosmic Ray Shielding Requirements During Solar Minimum*. NASA TM-4167, 1990.
29. Adams, James H., Jr.: *Cosmic Ray Effects on Microelectronics, Part IV*. NRL Memo. Rep. 5901 (Revised), Naval Research Lab., Dec. 31, 1987.
30. Wilson, John W.; and Badavi, Francis F.: *A Study of the Generation of Linear Energy Transfer Spectra for Space Radiations*. NASA TM-4410, 1992.
31. *1990 Recommendations of the International Commission on Radiological Protection*. ICRP Publ. 60, Pergamon Press Inc., c.1991.



(a) Projectile energy = 150 MeV/amu.




(b) Projectile energy = 600 MeV/amu.




(c) Projectile energy = 2400 MeV/amu.


Figure 1. Semiempirical, differential fragmentation cross section for liquid hydrogen target as function of projectile-ion charge Z_k and fragment-ion charge Z_j according to Rudstam's formalism for fragments heavier than helium. Cross sections shown for nucleon and helium fragments have been reduced by factors of 10 and 4, respectively. ($1 \text{ cm}^2/\text{g}$ converts to 1.67 barns.)



(a) Projectile energy = 150 MeV/amu.




(b) Projectile energy = 600 MeV/amu.




(c) Projectile energy = 2400 MeV/amu.


Figure 2. Semiempirical, differential fragmentation cross section for liquid hydrogen target as function of projectile-ion charge and fragment-ion charge according to Silberberg-Tsao formalism for fragments heavier than helium. Cross sections shown for nucleon and helium fragments have been reduced by factors of 10 and 4, respectively. (1 cm²/g converts to 1.67 barns.)



(a) Projectile energy = 150 MeV/amu.




(b) Projectile energy = 600 MeV/amu.




(c) Projectile energy = 2400 MeV/amu.


Figure 3. Semiempirical, differential fragmentation cross section for water target as function of projectile-ion charge and fragment-ion charge according to Rudstam's formalism for fragments heavier than helium. Cross sections shown for nucleon and helium targets have been reduced by factors of 10 and 4, respectively. (1 cm²/g converts to 29.9 barns.)



(a) Projectile energy = 150 MeV/amu.




(b) Projectile energy = 600 MeV/amu.




(c) Projectile energy = 2400 MeV/amu.

Figure 4. Semiempirical, differential fragmentation cross section for water target as function of projectile-ion charge and fragment-ion charge according to Silberberg-Tsao formalism for fragments heavier than helium. Cross sections shown for nucleon and helium fragments have been reduced by factors of 10 and 4, respectively. ($1 \text{ cm}^2/\text{g}$ converts to 29.9 barns.)




(a) 2-g/cm² thickness.




(b) 5-g/cm² thickness.

Figure 5. Differential LET spectra for annual dose equivalent behind various thicknesses of liquid hydrogen shield exposed to GCR at solar minimum.




(c) 15-g/cm² thickness.

Figure 5. Concluded.




(a) Differential LET spectra.




(b) Integral LET spectra.

Figure 6. Comparison of LET spectra for transmitted GCR flux calculated by using Silberberg-Tsao energy-dependent cross sections relative to the flux by Rudstam through several thicknesses of liquid hydrogen shield at solar minimum.




(a) Differential LET spectra.




(b) Integral LET spectra.

Figure 7. Comparison of LET spectra for transmitted GCR flux calculated by using Silberberg-Tsao energy-dependent cross sections relative to the flux by energy-independent cross sections through several thicknesses of liquid hydrogen shield at solar minimum.



(a) Differential LET spectra.



(b) Integral LET spectra.

Figure 8. Comparison of LET spectra for transmitted GCR flux calculated by using Silberberg-Tsao energy-dependent cross sections relative to the flux by Rudstam energy-independent cross sections through several thicknesses of liquid hydrogen shield at solar minimum.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY(Leave blank)	2. REPORT DATE August 1993	3. REPORT TYPE AND DATES COVERED Technical Paper		
4. TITLE AND SUBTITLE Semiempirical Fragmentation Models on Galactic Cosmic Ray Transport Calculations With Hydrogen Target		5. FUNDING NUMBERS WU 199-45-16-11		
6. AUTHOR(S) Judy L. Shinn, John W. Wilson, and Francis F. Badavi				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NASA Langley Research Center Hampton, VA 23681-0001		8. PERFORMING ORGANIZATION REPORT NUMBER L-17227		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA TP-3350		
11. SUPPLEMENTARY NOTES Shinn and Wilson: Langley Research Center, Hampton, VA; Badavi: Christopher Newport University, Newport News, VA.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT Unclassified-Unlimited Subject Category 93		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) Nuclear fragmentation cross sections of Silberberg and Tsao that are more accurate for a hydrogen target have been implemented in the data base to replace those of Rudstam for a galactic cosmic ray transport code (HZETRN). Sample calculations have been made for the transported galactic cosmic ray flux through a liquid hydrogen shield at solar minimum condition to determine the effect of such a change. The transported flux based on the Silberberg-Tsao semiempirical formalism contains fewer high-LET (linear energy transfer) components but more low-LET components than the results based on Rudstam's formalism; and this disparity deepens as the shield thickness increases. A comparison of the results obtained from using both energy-dependent and energy-independent cross sections of Silberberg and Tsao indicates that the energy-independent assumption results in an underestimation of high-LET flux above 100 keV/ μm by approximately 40 percent for a 15-g/cm ² thickness of liquid hydrogen. Similar results were obtained in a previous study (NASA TP-3243) when both energy-dependent and energy-independent cross sections of Rudstam were considered. Nonetheless, the present study found that an energy-independent calculation would be best accomplished by using Rudstam's cross sections as done in the past for various engineering applications.				
14. SUBJECT TERMS Galactic cosmic ray transport; Semiempirical nuclear fragmentation model		15. NUMBER OF PAGES 16		
		16. PRICE CODE A03		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

